

On random symmetric matrices with a constraint: the spectral density of random impedance networks

J. Ståring and B. Mehlig

School of Physics and Engineering Physics, Gothenburg University/Chalmers, SE-41296 Göteborg, Sweden

Yan V. Fyodorov

*Department of Mathematical Sciences, Brunel University, Uxbridge UB83PH, UK and
Petersburg Nuclear Physics Institute, Gatchina 188350, Russia*

J.M. Luck

*Service de Physique Théorique (URA 2306 of CNRS),
CEA Saclay, 91191 Gif-sur-Yvette cedex, France*

We derive the mean eigenvalue density for symmetric Gaussian random $N \times N$ matrices in the limit of large N , with a constraint implying that the row sum of matrix elements should vanish. The result is shown to be equivalent to a result found recently for the average density of resonances in random impedance networks [Y.V. Fyodorov, J. Phys. A: Math. Gen. **32**, 7429 (1999)]. In the case of banded matrices, the analytical results are compared with those extracted from the numerical solution of Kirchhoff equations for quasi one-dimensional random impedance networks.

The study of random matrices¹ has provided insight into many physical problems, both in the quantum and in the classical domain. For example, random matrices have been very successfully used to model statistical properties of disordered conductors and of highly excited classically chaotic quantum systems². In the classical domain random matrices arise, for instance, in the context of diffusion in random, directed environments (see for example Ref. 3 and references therein). In most of those applications the random matrix elements obeyed symmetry requirements where appropriate, and were otherwise taken to be independently distributed random variables.

There are cases, however, where constraints on the matrix elements must be considered. Such constraints generate correlations between the latter: examples are electron hopping in amorphous semiconductors (see Ref. 4 and references cited therein), random impedance networks^{5,6,7} (see Ref. 8 for a review and Ref. 9 for applications), and random master equations¹⁰: in these cases, the random matrices obey the constraints that the row sums of matrix elements should be zero. This condition implies correlations between diagonal and off-diagonal matrix elements. In Refs. 4,6,10, the average spectral density and spectral fluctuations of three different random-matrix ensembles of this type were calculated, using the method of replicas and the supersymmetry approach.

In this paper we derive the average density of eigenvalues of a suitably modified ensemble of symmetric Gaussian random matrices. The aim is twofold. First, we wish to show that for full matrices the density obtained is equivalent to that found in Refs. 6 and 10. Our second aim is to derive the corresponding result for banded matrices, and to compare it with the results of the numerical solution of the Kirchhoff equations on random impedance networks with quasi one-dimensional topology.

Formulation of the problem. We consider an ensemble

of $N \times N$ random matrices $\mathbf{M}$ with matrix elements

$$M_{mn} = J_{mn} - \delta_{mn} \sum_{l=1}^N J_{ml}, \quad (1)$$

where $\mathbf{J}$ is a real symmetric $N \times N$ matrix with random entries. Such ensembles have been considered in Refs. 6,7,10, and 11. In Ref. 10, $\mathbf{M}$ was used to model a random transition matrix for a model describing glassy relaxation, $\partial_t u(t) = -\mathbf{M}u(t)$, with matrix elements J_{mn} distributed independently (subject to the constraint $J_{mn} = J_{nm}$) according to

$$P(J_{mn}) = \frac{p}{N} \delta\left(J_{mn} - \frac{1}{p}\right) + \left(1 - \frac{p}{N}\right) \delta(J_{mn}). \quad (2)$$

The form (1) yields $\sum_n M_{mn} = 0$, implying probability conservation in the problem considered in Ref. 10. The average density $d(\lambda)$ of eigenvalues λ ,

$$d(\lambda) = N^{-1} \langle \text{tr} \delta(\mathbf{M} - \lambda \mathbf{1}) \rangle, \quad (3)$$

was calculated in the limit of large N and p , using the method of replicas. $\langle \cdots \rangle$ is an average over the ensemble defined by $P(J_{mn})$.

In Ref. 6, eigenvalues of matrices $\mathbf{M}$ with elements similar to (1) were shown to model resonance frequencies in random impedance networks⁵, with $J_{mn} = J_{nm}$ and

$$P(J_{mn}) = \frac{1}{2} \delta(J_{mn} - 1) + \frac{1}{2} \delta(J_{mn} + 1). \quad (4)$$

In Refs. 6 and 7, the average density of resonance frequencies was calculated, in the limit of large N , using a variant of the supersymmetry technique. It was found that the result agrees¹² with that of Ref. 10, up to a scale factor (related to p) and a rigid shift in λ [related to the fact that $J_{mn} \geq 0$ in (2)].




FIG. 3: Density of eigenvalues of exact diagonalisations of random matrices of the form (1,5) for $\sigma = 1$, $N = 500, 1000$, and $b = 50, 100$ (symbols), together with the prediction (13) (solid line).

where

$$g(z) = (2\pi)^{-1/2} \int_{-\infty}^{\infty} dJ \frac{\exp(-J^2/2)}{z - J}. \quad (9)$$

The result for the resolvent scales with σ^2 and N as $(\sigma^2 N)^{-1/2} g(H^{-1} (\sigma^2 N)^{1/2})$. Consequently, the equations shown in Fig. 2(a,b) imply the following self-consistent equation

$$G = (\sigma^2 N)^{-1/2} g((E - G) (\sigma^2 N)^{1/2}), \quad (10)$$

equivalent to Eq. (51) in Ref. 6 for the average density of resonances in an ensemble of highly connected random impedance networks¹⁵.

The form of the simplified contraction (8) implies an interpretation of the result (10) in terms of an averaged Pastur equation: consider a random matrix $\mathbf{M} = \mathbf{J} + \mathbf{V}$, where $\mathbf{J}$ is distributed according to (5), and $\mathbf{V}$ is a diagonal matrix with Gaussian random entries v_k with zero mean and unit variance, independent of J_{mn} . For a given realisation of $\mathbf{V}$, Pastur's equation¹⁴ is $\mathbf{G} = (E\mathbf{1} - \mathbf{V} - \mathbf{G})^{-1}$ (see also Ref. 16 and references therein). In this equation $\mathbf{G}$ is the $\mathbf{J}$ -averaged resolvent, keeping $\mathbf{V}$ fixed. One obtains Eq. (10) after observing that $\mathbf{G}$ is diagonal, by averaging over the matrix elements of $\mathbf{V}$. This interpretation implies that, in the limit of large N , the correlations between diagonal and off-diagonal matrix elements of $\mathbf{M}$ [as seen in Eqs. (1,7)] are irrelevant.

The above procedure is easily extended to the case of banded matrices, also of interest in random impedance networks^{5,11}. In the banded case, $\langle J_{mn} J_{kl} \rangle = \sigma^2 (\delta_{mk} \delta_{nl} + \delta_{ml} \delta_{nk})$ [which follows from (5)] is replaced by

$$\langle J_{mn} J_{kl} \rangle = \sigma^2 (|m - n|) (\delta_{mk} \delta_{nl} + \delta_{ml} \delta_{nk}). \quad (11)$$

The function $\sigma^2(x)$ is given by

$$\sigma^2(x) = \begin{cases} \sigma^2 & \text{for } 0 \leq x \leq b/2, \\ 0 & \text{otherwise.} \end{cases} \quad (12)$$




FIG. 4: Topology of a (quasi) one-dimensional lattice with periodic boundary conditions. Each site n is connected to its $b = 4$ neighbours.

The bandwidth of $\mathbf{J}$ is thus b . In the limit of large N and large b , the spectral density is given by a slight modification of (10),

$$G = (\sigma^2 b)^{-1/2} g((E - G) (\sigma^2 b)^{1/2}). \quad (13)$$

Diagrammatically, the necessary changes are most easily derived by letting $\sigma^2 = N^{-1}$ and assuming that $b = BN$ (with fixed $B \ll 1$). Then the wavy line in Fig. 1(a) acquires a factor of B . Furthermore, the dashed line in Eq. (8) also acquires a factor of B . Hence the self-consistency equation in the banded case becomes (13). Let us also note that the same formula is actually valid not only for $b \sim N$, but more generally for $1 \ll b \ll N$.

This result implies that densities for different values of b can be scaled on one single curve by plotting $\sqrt{b} d(E/\sqrt{b})$. In Fig. 3, solutions of (13) for $\sigma = 1$ are compared with results of exact diagonalisations of random matrices with $N = 500, 1000$ and $b = 50, 100$. We observe a very good agreement. The results confirm that, for large N and b , the average density of eigenvalues is independent of N , and that it scales with b as expected.

In the following we show that (13) also describes the density of resonances for certain random impedance networks with a large, but finite connectivity.

Random impedance networks. Random networks of complex impedances are currently used to model electrical and optical properties of disordered inhomogeneous media⁸. The most common situation is that of a binary composite medium, modeled by attributing a random conductance to each bond $(\mathbf{x}, \mathbf{y})$ of a lattice, according to the binary law:

$$\sigma_{\mathbf{x}, \mathbf{y}} = \begin{cases} \sigma_0 & \text{with probability } p, \\ \sigma_1 & \text{with probability } q = 1 - p. \end{cases} \quad (14)$$

The homogeneous Kirchhoff equations for the electric po-




FIG. 5: Plot of the density of resonances for a quasi one-dimensional random impedance network (see text). Data for three different ranges ($b = 60, 100$, and 120) (symbols) are compared with the theoretical prediction (13).

tentials,

$$\sum_{\mathbf{y}} \sigma_{\mathbf{x},\mathbf{y}} (V_{\mathbf{y}} - V_{\mathbf{x}}) = 0, \quad (15)$$

can be recast as

$$(\Delta_Q - \lambda \Delta)V = 0, \quad (16)$$

with $\lambda = \sigma_0/(\sigma_0 - \sigma_1)$, and

$$(\Delta V)_{\mathbf{x}} = \sum_{\mathbf{y}(\mathbf{x})} (V_{\mathbf{y}} - V_{\mathbf{x}}),$$

$$(\Delta_P V)_{\mathbf{x}} = \sum_{\mathbf{y} \in P(\mathbf{x})} (V_{\mathbf{y}} - V_{\mathbf{x}}), \quad (17)$$

$$(\Delta_Q V)_{\mathbf{x}} = \sum_{\mathbf{y} \in Q(\mathbf{x})} (V_{\mathbf{y}} - V_{\mathbf{x}}),$$

where $\mathbf{y}(\mathbf{x})$ are all the sites connected to site $\mathbf{x}$, whereas $\mathbf{y} \in P(\mathbf{x})$ (resp. $\mathbf{y} \in Q(\mathbf{x})$) are those connected by a conductance σ_0 (resp. σ_1), so that $\Delta = \Delta_P + \Delta_Q$. Resonances appear as non-trivial solutions to Eq. (16), for $0 < \lambda < 1$.

An efficient algorithm allowing for an exact determination of all the resonances of a finite sample has been developed in Ref. [5]. We have adapted this algorithm to the simplest geometry allowing for long-ranged bonds. Each site n of a very long chain is connected to its b neighbors ($n - b/2, \dots, n - 1, n + 1, \dots, n + b/2$), as shown in Fig. 4. Periodic boundary conditions are assumed. For definiteness we choose $p = q = 1/2$, so that all the resonances are expected to be located at $\lambda = 1/2$ in the $b \rightarrow \infty$ limit.

Our numerical results are shown in Fig. 5, for very long periodic chains with ranges $b = 60, 100$, and 120 . For each value of b , we have accumulated a number of resonances of order 10^7 . After rescaling the resonances according to $\lambda = (1 + E)/2$ the density is given by (13) with $\sigma = 1$. A very satisfactory quantitative agreement with the theoretical prediction is observed.

Acknowledgements. B.M. gratefully acknowledges discussions with J.T. Chalker. This work was in part supported by Vetenskapsrådet (J.S. and B.M.), and by EP-SRC Research Grant No. GR/13838/01 (Y.V.F.).

- ¹ M.L. Mehta, *Random Matrices and the Statistical Theory of Energy Levels* (Academic Press, New York, 1991).
- ² Th. Guhr, A. Müller-Groeling, and H.A. Weidenmüller, *Phys. Rep.* **299**, 189 (1998).
- ³ J.T. Chalker and Z.J. Wang, *Phys. Rev. Lett.* **79**, 1797 (1997); *Phys. Rev. E* **61**, 196 (2000).
- ⁴ M. Mézard, G. Parisi, and A. Zee, *Nucl. Phys. B* **559**, 689 (1999); C.R. Offer and B.D. Simons, *J. Phys. A: Math. Gen.* **33**, 7567 (2000).
- ⁵ Th. Jonckheere and J.M. Luck, *J. Phys. A: Math. Gen.* **31**, 3687 (1998).
- ⁶ Y.V. Fyodorov, *J. Phys. A: Math. Gen.* **32**, 7429 (1999).
- ⁷ Y.V. Fyodorov, *Physica E* **9**, 609 (2001).
- ⁸ J.P. Clerc, G. Giraud, J.M. Laugier, and J.M. Luck, *Adv. Phys.* **39**, 191 (1990).
- ⁹ F. Brouers et al., *Physica A* **241**, 146 (1997); *Phys. Rev. B* **55**, 13234 (1997); E.M. Baskin et al., *Physica A* **242**, 49 (1997); M.V. Entin, *Zh. Eksp. Teor. Fiz.* **114**, 669 (1998); S. Grésillon et al., *Phys. Rev. Lett.* **82**, 4520 (1999); A.K. Sarychev and V.M. Shalaev, *Physica A* **266**, 115 (1999); A.K. Sarychev and V.M. Shalaev, *Phys. Rep.* **335**, 275 (2000); L. Zekri et al., preprint cond-mat/9908487.

- ¹⁰ A.J. Bray and G.J. Rodgers, *Phys. Rev. B* **38**, 11461 (1988).
- ¹¹ Y.V. Fyodorov, *JETP Letters* **70**, 743 (1999).
- ¹² The average density of resonance frequencies of an ensemble of random impedance networks is obtained from a generalised eigenvalue problem (random matrix pencil) of the form $\mathbf{M}u = \lambda \mathbf{W}u$, with $W_{mn} = (N + 2)\delta_{mn} - 1$. It was shown^{6,7} that, as a consequence of the particular structure [see eq. (1)] of $\mathbf{M}$, and of the form of $\mathbf{W}$, the latter can be replaced by N times the unit matrix without changing the average density of eigenvalues.
- ¹³ The constant of proportionality is fixed by normalisation. To compare with Refs. 5 and 8, the variance should be set unity; for the perturbative calculations described below it is convenient to take $\sigma^2 = N^{-1}$.
- ¹⁴ L.A. Pastur, *Theor. Math. Phys. (USSR)* **10**, 67 (1972).
- ¹⁵ Note the missing factor 1/2 in the right-hand side of Eq. (51) of Ref. 6.
- ¹⁶ Y.V. Fyodorov and A.D. Mirlin, *Phys. Rev. B* **52**, R11580 (1995).